

EVALUATION OF ENSILING QUALITIES OF SIGNAL GRASS (*Brachiaria mulato II*) AND COOK STYLO (*Stylosanthes guianensis*) AT DIFFERENT PROPORTION

^{1*}GAMBO, S., ²MUHAMMAD, I. R., ¹ABDULLAHI, A. Y., ¹IBRAHIM, U., AND
³MUHAMMAD, A. S.

¹Department of Animal Science, Aliko Dangote University of Science and Technology, PMB
3244, Wudil, Kano State, Nigeria.

²Federal University of Agriculture, Zuru, PMB 28, Zuru Kebbi State, Nigeria.

³Department of Animal Health and Production, Binyaminu Usman Polytechnic, PMB 013,
Hadejia Jigawa State, Nigeria.

*Corresponding Author: sulaimangambo89@gmail.com (+2347082274742)

Abstract

The study aimed to evaluate the ensiling qualities of *Stylosanthes guianensis* (Cook), grown with *Brachiaria mulato II* (Signal grass) at different mixing ratio. The experiment was conducted at Animal Science Laboratory of the Faculty of Agriculture, Bayero University Kano. The treatments were the mixture of *Stylosanthes guianensis* and *Brachiaria mulato II* at 100:0, 75:25, 50:50, 25:75 and 0:100% ratios laid in a completely randomized design (CRD) and replicated three thrice. Fresh *Brachiaria mulato II* and *Stylosanthes guianensis* were harvested, wilted for 24 hours and chopped to 2-3cm long for ease of compaction and combination for silage. The research materials in the silos were sealed tightly and allowed for 28 days fermentation period. At expiration, the resultant silages were scored for colour and aroma by three independent scorers. The results of silage study showed that irrespective of different mixing proportions of *Stylosanthes guianensis* – *Brachiaria mulato II*, the colour of the ensiled mixtures were yellowish green with pleasant and very sweet aroma. The pH recorded was slightly acidic in all the treatment combinations with a temperature range of 24.6^oC – 29.6^oC. The crude protein (CP) and ether extract content of the silage were not significantly ($p>0.05$) different, but numerically higher than the mean value of the former obtained in *Stylosanthes guianensis* (50%) *Brachiaria mulato II* (50%). The study concluded that the use of legume in preparation of grass silages has advantages to silage quality characteristics and nutritional composition. The result obtained revealed that the composite silage made from 50% Mulato II plus 50% stylo resulted in silage with higher CP and a better dry matter intake content.

Keywords: Cook stylo, signal grass, silage, nutrients

Introduction

The major constraint to livestock production in developing countries such as Nigeria is the scarcity and fluctuating quantity and quality of all year-round feed supply (Kallah, *et al.*, 1997). Forages are essential for the successful operation and a highly profitable animal production venture. This is more relevant to ruminants which rely heavily on forages for their health and production in a cost effective and sustainable manner (Addo, 2019). Forage and fodder crops are essential in the development of national livestock industry as cultivated fodder crops and other feed resources partake heavily in the contemporary animal agriculture (Rusdy, 2016). About 50-80% of Nigerians are involved in crop, livestock or crop-livestock agriculture. The quality and quantity of feed resources are the major constraints of increasing ruminant productivity under tropical set up (Addo, 2019). Grasses due to their adaptations are advantageous over other plant species as they have surface rooting system and can respond effectively to management (February and Haggins, 2010). Grasses generally are typically known to have low crude protein

that cannot solely sustain ruminant animals throughout the year, hence the need for intercropping with forage legumes. One way of increasing the grazing resources of natural pasture is to integrate forage legumes into the pastures, with the aim of diversifying the sources of forage and at the same time increasing the amount of protein available for the grazing animals as well as increasing the nitrogen uptake of associated forage grass (Macharia, 2003). The importance of forage legumes in increasing herbage production from grasses and to enhance the quality of feed produced has been recognized (Muinga *et al.*, 2007).

The seasonal variation in weather and climate results in feed inadequacy in quantity and quality in the tropical region. An alternative for alleviating this challenge is through ensiling the excess forage produced during the rainy season. It permits forage storage in amounts and quality to be used for animal feeding during the dry season, and serves as an alternative for roughage supplementation during dry season (Balsalobre *et al.*, 2001). However, Tropical forages are susceptible to losses during ensilage due to their low buffering capacity, humidity higher than 70% and low water - soluble carbohydrate (WSC) content (Balsalobre *et al.*, 2001). Silage can be defined as any plant material that has undergone fermentation or “pickling” in a silo. Silo is any storage structure in which green, moist forage is preserved (Stewart, 2011). The primary goal of making silage is to maximize the preservation of original nutrients in the forage crop for feeding livestock at a later date in livestock feeding programs (Stewart, 2011). The first essential objective is to achieve anaerobic conditions under which natural fermentation can take place. In practice this is achieved by consolidating and compacting the material and the sealing of the silo to prevent re-entry of air. Air that is trapped in the herbage is rapidly removed by respiratory enzymes (McDonald *et al.*, 1991). Where oxygen is in contact with herbage for a period of time, aerobic microbial activity occurs and yeast would grow. This causes the material to decay to a useless, inedible and frequently toxic product (McDonald *et al.*, 1991). Finer chopping of plant material results in improved compaction and fermentation of silage. This then improves palatability and intake of silage (Meeske, 2005).

The preservation of excess pasture during surplus in the form of silage is indispensable and this allows mitigating the feed deficit that could be encountered during the dry season. Ensiling is a potent general method for forage preservation and also a form of treatment to occasionally salvage the under-utilized pastures for better acceptability and degradability. The quality of silage is however dependent on the quality of the crop at ensiling, type of fermentation, rate of pH decrease, moisture content of the crop and anaerobic conditions (Meeske *et al.*, 2000). Silage making in the tropics is paramount if there will be an all year-round availability of forages for livestock. Fodder conservation is promoted with the main objective of ensuring feed availability during periods of feed limitation (Moh'd *et al.*, 1993). Silage quality depends on many factors such as plant species, implemented agro-technical measures and practices, preparation time, applied inoculants, especially stages of plant development in which it is used for the preparation of silage. McEniry *et al.* (2013) stated that the two most important pre-ensiling factors responsible for the quality of silage are plant species and stage of plant development at the time of harvest. The study evaluated the ensiling qualities of *Stylosanthes guianensis* (Cook), grown with *Brachiaria mulato* II (Signal grass) at different mixing ratios.

Materials and Methods

Experimental Location, Design and Treatment

The experiment was conducted at the Animal Science Laboratory of the Faculty of Agriculture, Bayero University Kano. The experimental area is located on latitude 11.97095°N of the equator and longitude 8.42511°E of the Greenwich meridian' (GPS Coordinate, 2019). The treatments were mixtures of *Stylosanthes guianensis* and *Brachiaria mulato* II at 100:0, 75:25, 50:50, 25:75 and 0:100% ratios laid in a completely randomized design (CRD) and replicated three times.

Silage Making

Silage was made in a laboratory silo (500 gram capacity bottle). Fresh *Brachiaria mulato* II and *Stylosanthes guianensis* were harvested from the field and allowed wilted for 24 hours in order to reduce the moisture content after which were chopped to 2-3cm long for ease of compaction and combination for silage. The materials in the silos were sealed tightly and allowed for 28 days fermentation period. At expiration, the resultant silages were scored for colour and aroma by three independent scorers on a scale of 1 - 4 as shown in Table 1, Thereafter pH and temperature were determined using the digital pH meter and thermometer respectively.

Table 1: Description of colour and aroma rating used as indices of silage quality

Scale	Colour	Aroma
1	Dark/Deep brown	Putrid or rancid
2	Light brown	Pleasant
3	Pale yellow	Sweet
4	Yellowish green	Very sweet

Source: Muhammad *et al.* (2009).

Chemical Analysis

Silage samples collected were oven dried according to the treatments and milled to pass through 2mm sieve and stored in a laboratory container. The milled samples were subjected to proximate analysis to determine crude protein (CP), Ash, crude fibre, ether extract (fat) and nitrogen free extract according to procedures described by AOAC (1995). Neutral Detergent Fibre (NDF) and Acid Detergent Fibre (ADF) were determined using the method of Van Soest *et al.* (1991).

Statistical Analysis

Data obtained in this study were analyzed using statistical analysis system (GENSTAT, 2014) Version 17.1. Significant means were separated using Least Significant Difference (LSD) at 5% level of probability.

Results and Discussion

Silage Characteristics

The results of pH, temperature, aroma and colour characteristics of silage made at different grass and legume mixing proportions are presented in Table 2. The mean values of pH obtained in Mulato II (75%) – Cook stylo (25%) (T4) were significantly different ($p < 0.05$) from other treatment combinations. All the silage samples prepared were slightly acidic with pH value ranges from 5.2 - 5.8.

Table 2: Effect of grass-legume mixture at different proportion on silage characteristics

Treatment	pH	Temperature (⁰ C)	Aroma	Colour
Mulato II (0%) – Cook stylo (100%)	5.410 ^b	29.60 ^a	1	1
Mulato II (25%) – Cook stylo (75%)	5.213 ^b	28.57 ^a	2	4
Mulato II (50%) – Cook stylo (50%)	5.310 ^b	29.17 ^a	2	4
Mulato II (75%) – Cook stylo (25%)	5.827 ^a	26.07 ^b	4	4
Mulato II (100%) – Cook stylo (0%)	5.457 ^b	24.60 ^b	3	3
LSD	0.3068	2.402	-	-

Means within the same column with different superscripts differ significantly (p<0.05)

Effect of mixing proportion of *Brachiaria mulato II* with *Stylosanthes guianensis* on proximate composition and feeding value of silage

Effect of mixing proportion of *Brachiaria mulato II* with *Stylosanthes guianensis* on proximate composition and feeding value of silage is presented in Table 3. All the parameters evaluated were significantly different (p<0.05), except CP and EE. The numerically higher mean value of CP content (16.99 %) was obtained in T3 and decrease with the decrease in legume proportion. The relative feed value (RFV) in T3 (84.52%) was observed to be higher among the treatment combination and absolutely revealed dry matter intake (DMI) that is not significantly different from the treatments with higher proportion of legume in grass-legume mixtures.

The pH value is one of the simplest and quickest ways of evaluating silage quality; the lower the pH, the better the preserved and more stable is the silage. Silage that is properly fermented will have a much lower pH (more acidic to slightly acidic) than the original forage. The pH value of good silage below 4.1 has been considered as an index of good quality (Harrison *et al.*, 1995; Amuda *et al.*, 2020). The pH values of all the silage samples in this study were within the range value (5.5-6.1) of ensiled gamba grass fortified with tropical legumes (centrosema, lablab and groundnut leaves) Muhammad *et al.* (2009). The decreasing pattern of pH value with the increase in grass proportion in the mixtures is in agreement with the previous report by Kung *et al.* (2001) who stated that, the highest pH values were in treatments with the most favourable seeding ratios for legumes, and this might be attributed to the increased buffering power of the leguminous ensiled biomass. Temperature is one of the essential factors affecting silage colour. The lower the temperature during ensiling, the less will be the colour change as excessive heat owing to high temperature do affect silage colour (Adesogan and Newman, 2010). However, the range of temperature (24.6⁰-29.6⁰C) obtained during the fermentation period of 28 days was not in agreement with the temperature 26.0⁰-26.3⁰C reported by Amuda *et al.* (2017) for ensiled maize stover with and without additive.

The proximate composition of ensiled Mulato II, Stylo and their mixtures at different mixing proportions in all the parameters evaluated revealed significant differences except in CP and EE across the treatments. The least CP value was obtained in sole Mulato II (14.77%) which was higher than the value reported by Amuda *et al.* (2020) with 36.09%.

Table 3: Effect of mixing proportion of *Brachiaria mulato II* with *Stylosanthes guianensis* on proximate composition and other feed values of silage (%)

TRT	DM	MOIST	ASH	CF	CP	EE	NFE	ADF	NDF	DMI	DDM	RFV
T1	93.4 ^b	6.60 ^a	9.10 ^c	15.98 ^c	15.94	4.00	61.39 ^a	40.18 ^{cd}	68.58 ^b	1.750 ^a	57.60 ^{ab}	84.00 ^{ab}
T2	93.50 ^b	6.50 ^a	11.35 ^b	16.22 ^c	16.50	4.80	60.32 ^{ab}	40.69 ^c	70.36 ^b	1.706 ^a	57.20 ^b	81.30 ^b
T3	94.20 ^{ab}	5.81 ^{ab}	11.40 ^b	17.31 ^c	16.99	4.85	57.02 ^{bc}	38.86 ^d	69.37 ^b	1.730 ^a	58.63 ^a	84.52 ^a
T4	93.48 ^b	6.53 ^a	9.38 ^c	27.00 ^a	14.87	4.91	47.33 ^d	45.85 ^a	75.74 ^a	1.585 ^b	53.18 ^d	70.22 ^d
T5	94.58 ^a	5.43 ^b	12.53 ^a	22.43 ^b	14.77	3.40	54.22 ^c	43.86 ^b	73.92 ^a	1.624 ^b	54.73 ^c	70.05 ^c
P- value	0.048	0.048	0.003	0.001	0.644	0.033	0.001	<.001	0.002	0.002	<.001	<.001
SEM	0.2066	0.2066	0.272	0.388	1.193	0.233	0.855	0.360	0.491	0.0111	0.280	0.787

Means within the same column with different superscripts differ significantly ($p < 0.05$). DM=Dry matter, MOIST. =Moisture, CF= Crude fibre, CP= Crude protein, EE= Ether extract, NFE= Nitrogen free extract, ADF= Acid detergent fibre, NDF= Neutral detergent fibre, DMI= Dry matter intake, DDM= Digestible dry matter, RFV= relative feed value

The CP content of silages ranged from 9.77 – 16.99% and higher than the range 9.44 - 14.88% recorded by Amuda *et al* (2020) in gamba grass ensiled with different graded levels of *Centrosema pubescence* and was contrarily to the findings that, the higher CP value are obtained from high levels of legume inclusion Kallah *et al.* (1997). Silage made with Mulato II Stylo at 50:50 (T3) revealed a higher CP content (16.99%) and the result was in line with the fact that the use of Stylo and Centro improved the silage quality of grass (Muhammad *et al.*, 2008). In contrast, the higher the level of legume inclusion; the higher the CP and conversely, the lower the CF content of the silage (Muhammad *et al.*, 2009). The CP content observed in this study in all the treatments was within the NRC (2001) recommended percentage of 12% and 16.5% as CP required for young and lactating animals respectively. However, the CP in silages increased after ensiling, and mixed silages had higher CP than sole silages. Although, according to an earlier study, CP decreases after ensiling because of oxygen consumed during respiration and proteolysis at the onset of ensiling (McDonald *et al.*, 1991). Another recent study showed similar results with the findings of this study that the increase in CP is due to excess protein produced by the fermentative microbial communities (Lai *et al.*, 2022). The higher CF content (27.00%) in T4 (Stylo:Mulato II) (25:75) and lower in sole Stylo observed in this study was similar to the findings of Moran (2005) and Elkholy *et al.* (2009); but lower than the least values of 33.10 and 36.30 % reported by Ajayi *et al.* (2012) and Amuda *et al.* (2020), respectively. Ash content was not within the favourable ash content of grass and legume forages of 7 – 9%, but is significant to what was reported by Ajayi *et al.* (2012) and Amuda *et al.* (2020), respectively.

Nitrogen free extract indicates the soluble carbohydrate of the forage plant. The range (47.33 – 61.39%) obtained was higher than the range (35 – 53%) in ensiled gamba grass with graded levels of centro and obtained in maize stover silages (Amuda *et al.*, 2020). Sanderson; (2010) reported that the proportion of grasses is positively correlated with the fibre and digestibility of legume–grass mixtures (Sturludóttir *et al.*, 2014). Sing and Oosting (1992) reported that roughage feeds containing values of less than 45% could be classified as high quality, those with values ranging from 45-65% as medium with values and those with higher than 65% as low quality. The RFV is used to predict the intake and energy content of the forages and the higher it is the better the nutritional value of the forage. The value obtained in this study was positively correlated with the CP content, suggesting that the treatment combination with higher value revealed a higher RFV and this might be attributed to the contribution of legume proportion.

Conclusion

This study has shown that mixture proportions have significant effect on the dry matter yield and nutritional quality of legume–grass mixtures rather than species diversity. It can be concluded that the use of legumes in preparation of grass silages has advantages to silage quality characteristics and nutritional composition. This study has revealed that the composite silage made from 50% Mulato II plus 50% stylo resulted in silage with higher CP and a better dry matter content.

References

- AOAC (1995). Association of Official Analytical Chemists Official Methods of Analysis (15th edition), Washington D.C., USA Pp. 69-88.
- Addo, K. A. (2019). Propagation, forage production and forage quality of some Ghanaian browse plants, University of Ghana: Digital collection. <http://197.255.68.203/handle/123456789/6807>.
- Adesogan, A. T. & Newman Y. C. (2010). Silage harvesting and feeding SS-AGR-177/AG180-*Forage Management (Florida Forage Handbook)*, Pp.1-7.
- Ajayi, F. T. (2012). Effects of feeding ensiled mixture of elephant grass (*Pennisetum purpureum*) with three grain legume plants on digestibility and nitrogen balance of West African Dwarf Goats. *Livestock Science*.142, (1); 80 – 84.
- Amuda, A. J., Falola, O. O. & Babayemi, O. J. (2017). Chemical composition of quality characteristics of ensiled maize stovers. Federal University Wukari *Trends in Science and Technology Journal*, 2 (1), 195-198.
- Amuda, A. J., Alabi, B.O. & Jonah, S. A. (2020). Nutritional quality of ensiled gamba grass (*Andropogon gayanus*) fortified with graded levels of Centro (*Centrosema pascourum*). *Nigerian Journal of Animal Production*, 47(1); 234 – 245.
- Balsalobre, M. A. A., Nussio, L. G. & Martha, J. G. B. (2001). Controle de Perdasnaproducao de Silagens de Gramineastropicais. In A Pruducacao Animal na Visao dos Brasileiros Piracicaba, Brazil: FEALQ. Retrieved from <http://sbz.org.br/documentos--anais-das-reunioes>.
- Elkholy, M. E. H., Hassanein, E. I. I., Soliman, M. H., Eleraky, W. E., Elgameiel, M. F. A. & Ibraheim, D. (2009). Efficacy of feeding ensiled corn crop residues to sheep. *Pakistan Journal of Nutrition* 8 (12); 1856 –1867.
- February, E. C. & Higgin, K. (2010). The distribution of tree and grass roots in Savannas in relation to soil nitrogen and water.; *South African Journal of Botany*; 76, 517.
- GENSTAT, (2014). General Statistics Software for Windows 15th Edition. VSN International, Hemel Hempstead, UK. Web page: Genstat.co.uk.
- Harrison, J. H., Blauwiekel, M. R. & Stokes, M. R. (1995). Fermentation and utilization of grass silage. *Journal of Dairy Science*; 77; 3209–3235 <https://www.environment.gov.au/system/files/pages/99dfad7e-9feb-4da1-826b>.
- Kallah, M. S., Baba, M., Alawa, J. P., Muhammad, I. R. & Tanko, R. J. (1997). Ensiling quality of Columbus grass (*Sorghum almum*) grown in Northern Nigeria. *Animal Feed Science and Technology*, 68;153-163.
- Kung, L. & Shaver, R. (2001). Interpretation and use of silage fermentation analysis reports. *Focus on Forage*, 3(13); 1–5.
- Lai, H., Gao, F., Hao, S., Zheng, P., Li, Y. & Yao, H. (2022). Nitrogen distribution and soil microbial community characteristics in a legume–cereal intercropping system: A Review. *Agronomy* 12;1900. doi: 10.3390/agronomy12081900
- Macharia, P. N. (2003). Introduction of forage legumes into natural pastures of semi-arid rangelands of Kajiado District of Kenya. Kenya Agricultural Research Institute (KARI) *Legume Research Network Project Newsletter* Issue No. 10.
- McEniry, J., King, C. & O’Kiely, P. (2013). Silage Fermentation Characteristics of Three Common Grassland Species in Response to Advancing Stage of Maturity and Additive Application. *Grass and Forage Science*, 69, 393-404.

- McDonald, P., Henderson, A. R. & Heron, S. J. (1991). The Biochemistry of Silage. Chalcombe Publications, Aberystwyth, UK, 32-37.
- Meeske, R., Cruywagen, C.W., Vander Merwe., G. D. & Greyling J. F. (2000). The effect of adding lactic acid bacterial inoculant to a big round bale Oat silage on intake, milk production and milk composition of Jersey Cows. *South African Journal of Animal science*; 30; 80-81.
- Meeske, R. (2005). Silage additives: Do They Make a Difference? Vol. 6, Department of Agriculture Western Cape, Ellensburg, X1. *South Africa Journal of Animal Science*. (n.p.)
- Mohd, N. M. A., Aminah, A. & Idris A. B., (1993). Forage conservation for Livestock Smallholders in Malaysia. Pp. 103-109, In: Strategies for suitable forage based livestock production in Southeastern Asia. *Proceedings of the 3rd Meeting of the Regional Working Group on Grazing and Feed Resources of Southeast Asia, Hhonkaen, Thailand*.
- Moran, J.; (2005). Tropical Dairy Farming: Feeding Management for Smallholder Dairy Farmers in the Humid Tropics. Land link press. Pp. 321.
- Muhammad, I. R, Baba, M. Mustapha, A., Ahmed, M. Y & Audu-Rahman, L. S., (2008). Use of legumes in the improvement of silage quality of Columbus grass. *Medwell Research Journal of Animal Sciences* 2(4); 109-112.
- Muhammad, I. R., Abdu, M. I., Iyeghe-Eraktobor, G.T. & Sulaiman K. A. (2009). Ensiling quality of Gamba fortified with tropical legumes and its preference by rabbits. *Research Journal of Applied Sciences*, 4 (1); 20-25.
- Muinga, R.W., Mureithi, J. G., Juma, H. & Saha, H. M., (2007). “The effect of supplementing Napier grass or maize stover basal diet with Either *Gliciridia*, *Clitoria* or *Mucuna* on manure quality and quantity in Jersey Cows.” *Tropical and Subtropical Agroecosystems*, 7(3); 157-163.
- NRC (2001). National Research Council. *Nutrient Requirements for Dairy Cattle*, 7th Edn. National Academic of Sciences, Washington, D.C.
- Rusdy, M. (2016). Elephant grass as forage for ruminant animals. Department of Forage Crop and Grassland Management, Faculty of Animal Science, Hussanuddin University, Indonesia.
- Sanderson, M. A. (2010). Nutritive value and herbage accumulation rates of pastures sown to grass, legume, and chicory mixtures. *Agronomy Journal*, 102; 728–733.
- Singh, G. P. & Oosting, S. J. (1992). A model for describing the energy value of straws. *Indian Dairyman* XLIV; 32- 327.
- Stewart, W. M.; (2011). Plant Nutrition Today, *Scientific Journal of Applied Science*, 4; 5.
- Sturludóttir, E., Brophy, C., Belanger, G.; Gustavsson, A. M., Jørgensen, M., Lunnan, T. & Helgadóttir, Á. (2014). Benefits of mixing grasses and legumes for herbage yield and nutritive value in Northern Europe and Canada. *Grass Forage Science*, 69; 229–240.
- Van Soest, P. J., Robertson, J. B. & Lewis, B. A. (1991). Methods for dietary fibre, neutral detergent fibre and non-starch polysaccharides in relation to Animal Nutrition. *Journal of Dairy Science*, 74; 3583-3597.